


Hadronic Structure from Perturbative Dressing in QCD - The Valon Model

Firooz Arash^(a) and Fatemeh Taghavi^(b)

^(a) *Physics Department, Tafresh University, Tafresh, Iran*

^(b) *Physics Department, Iran University of Science and Technology, Narmak, Tehran, Iran*



In the Framework of *valon model* we have calculated the parton distribution in a valon for both the polarized and unpolarized cases. These distributions are originated purely from the dressing of a valence quark in QCD and are common to all hadrons. Structure functions, $F_2^{p,\pi}$, are obtained, which agree rather well with the experimental results. A simple relation between F_2^p and F_2^π is inferred. For the polarized structure function, g_1 while the model gives good agreement with data, it requires a sizable angular momentum contribution to the spin of the valon, and hence to that of proton. This contribution is calculated.

Introduction The parton distributions in proton have been studied extensively in a wide range of both x and Q^2 . The parton distribution functions (PDF) are calculated numerically by fitting over 1300 data points which rely on over 30 parameters in the input PDF's. Although these PDF's are accurate but inconvenient to describe analytically. Recently, some experimental results have also emerged for the pion¹, at very low x which can serve as a further check on our understanding of hadron structure. Here we elaborate on the valon model that can be very useful in the study of hadronic structure, in particular when the experimental data are scarce. A valon has its own cloud of partons which can be calculated in pQCD. This structure is universal and independent of the hosting hadron. In² the structure of the valon is calculated to the Next-to-leading Order in QCD and tested against the $F_2^p(x, Q^2)$ in a wide range of kinematics: $x = [10^{-4}, 1]$ and $Q^2 = [1, 5000] \text{ GeV}^2$. Here we will examine its ability in describing pion structure functions, F_2^π without any additional new parameter. This is important, because at first it is not clear at all that if one moves away from proton, where there are ample data points, to pion where the data are much restricted, the insight gained from the study of proton, will still be valid. At the end we will discuss the case of polarized structure function g_1 .

Formalism: Unpolarized The unpolarized structure of a valon is calculated in next-to-leading

^ae-mail: farash@cic.aut.ac.ir

order framework of PQCD in Ref.[2]. Without going into the details it suffice here to give the parameterized form of the valonic PDF as follows:

$$zq_{valence}^{valon}(z, Q^2) = az^b(1-z)^c, \quad (1)$$

$$zq_{sea}^{valon}(z, Q^2) = \alpha z^\beta(1-z)^\gamma[1 + \eta z + \xi z^{0.5}]. \quad (2)$$

The parameters $a, b, c, \alpha, \beta, \gamma, \eta$, and ξ are functions of Q^2 and are given in the appendix of Ref. [2]. Gluon distribution in a valon has an identical form as in Eq. (2) but with different parameter values. Upon convolution of these valonic PDF's with the valon distribution, $G_{valon}^h(y)$ in a hadron, we obtain the hadronic structure function:

$$F_2^p(x, Q^2) = \sum_{valon} \int_x^1 dy G_{valon}^h(y) F_2^{valon}(\frac{x}{y}, Q^2) \quad (3)$$

where $G_{valon}^h(y)$ is the distribution of valon with momentum fraction y in hadron h . These functions are given in Refs.^{2,3,4} for proton, pion, and kaon. F_2^{valon} is the structure function of a valon. In figures (9) of Ref.[2] the results for the proton structure function is presented at $1 < Q^2 < 2000 \text{ GeV}^2$. They are compared with the experimental values and with other global fitting results, the agreements are excellent. For the details see Ref.[2].

Pion structure function at low x in the leading neutron experiment [1] is reported. Moreover, there are some data from Drell-yan⁵ experiments at medium and large x for the valence quark distribution. We have used the data of Ref.[5] at $Q^2 = 25 \text{ GeV}^2$ to determine $G_{valon}^\pi(y)$. In Figure [1] the results are shown along with the SMRS and GRS determination. It turns out that the valon distribution in pion is a very broad:

$$G_{\pi^+}^{\frac{U}{\pi^+}}(y) = G_{\pi^+}^{\frac{\bar{D}}{\pi^+}}(y) = 1.02(1-y)^{0.01}y^{0.01} \quad (4)$$

This is expected; for it indicates that the valons in a pion are tightly bound, a reflection of the fact that the pion is much lighter than the mass of its valons. From Eqs.(1-5) F_2^π is readily calculated without any free parameters. In Figure (2) we present F_2^π at $Q^2 = 7$ and $Q^2 = 15 \text{ GeV}^2$ at very low x along with the data from Ref.[1] and the results from SMRS and GRS. In the figure there are two sets of data points, corresponding to two different normalization of the data by ZEUS coll. They differ by a factor of two [1]. Our results favor the effective flux normalization. Same is true for SMRS, but GRS is more compatible with the additive quark model normalization. However, as can be seen from figure (1) GRS determination grossly disagrees with the valence quark distribution in pion. In ³ and ⁴ it is shown that the valon description holds true for all values of Q^2 .

It is also interesting to notice that a simple relationship holds between F_2^p and effective flux normalization of pion structure function $F_2^{\pi(EF)}$ which is also noticed by ZEUS Coll. in Ref. [1]; namely:

$$F_2^{\pi(EF)}(x, Q^2) \approx k F_2^p(x, Q^2) \quad (5)$$

With $k \approx 0.37$. We have calculated both sides of equations (6) independently, using the valon model; a sample result at $Q^2 = 15 \text{ GeV}^2$ is given in Figure (3), but the relationships also hold true for other values of Q^2 .

Polarized Structure Functions The case of polarized structure function requires knowledge of polarized parton distribution in a polarized valon and the distribution of polarized valon in polarized hadron. We will restrict ourselves to nucleon. The first task is carried out similar to that of unpolarized case; namely, the moments of polarized structure functions in a valon is calculated in QCD, then with Inverse Mellin transform technique the polarized parton densities are obtained. As for the second requirement, we define the function, $\delta F_j(y)$ via:

$$\delta G_j(y) = \delta F_j(y) G_j(y) \quad (6)$$




Figure 1: **Right-** Comparison of the pion valence distribution, $u_{valence}^{\pi-}(x)$, using the valon model calculation (solid line), SMRS (dashed line), and GRS (dotted line), with the data from Ref.[1] at $Q^2 = 25 \text{ GeV}^2$. **Left-** Pion structure function, $F_2^\pi(x)$, at $Q^2 = 7 \text{ GeV}^2$. The diamonds and the squares are pion flux and additive quark model normalization of the data [1], respectively. The solid line is the model calculation; The dashed line is from SMRS and the dotted line is from GRS determination.




Figure 2: **Right** The same as the left figure in Figure-1 but at $Q^2 = 15 \text{ GeV}^2$. **Left-** The effective flux normalization of F_2^π data as compared with the scaled F_2^p at $Q^2 = 15 \text{ GeV}^2$ of equations (5). The data points are from Ref.[1]. The solid line is $0.371F_2^p(x, Q^2)$ pertinent to Eq. (5) and the dashed line is F_2^π , both are calculated directly and separately from the model.

where, j refers to U or D valon, and $G_J(y)$ is the unpolarized valon distribution. In addition, we impose upon $\delta F_j(y)$, the constraint that $P_U = 2/3$ and $P_D = -1/3$, corresponding to $SU(6)$ model, with P_j defined by $P_j = \int_0^1 dy \delta F_j(y) G_J(y)$. Then, for a nucleon we have:

$$g_1^N(x, Q^2) = \int_x^1 dy [2\Delta G_U g_1^U(\frac{x}{y}, Q^2) + \Delta G_D g_1^{UD}(\frac{x}{y}, Q^2)] \quad (7)$$

This completes the formalism. In Figure (4) xg_1^p are shown. It is evident that the model is able to reproduce the experimental data. Much interest has been centered on the integral

$$\Gamma^{p(n)} = \int dx g_1^{p(n)}(x). \quad (8)$$

Our model gives $\Gamma^p = 0.1212, 0.1219, 0.1225$, and 0.1236 for $Q^2 = 2, 3, 5, 10 \text{ GeV}^2$, respectively, which are very close to the experimental data. The same is true for Γ^n . These results suggest that quark contribution to the spin of nucleon is small. So, while the model produces all available data on polarized structure function, but the same data do not give the nucleon spin. Therefore, it is natural to ask if the spin content of a valon do give the spin $\frac{1}{2}$ of the valon itself. The contribution of different components of a valon to its spin is obtained by the following integral, say for a U-type valon.

$$\Delta H_{\frac{i}{U}}(Q^2) = \int_0^1 \delta q_{\frac{i}{U}}(z, Q^2) dz \quad (9)$$




Figure 3: **Right-** $xg_1^p(x, Q^2)$ as calculated from the model and compared with the experimental data. **Left-** various component contributions to the spin of a valon as a function of Q^2

where i is for valence, sea, and gluon inside a valon. For the valence part, $\Delta H_{\text{valence}}^U$ is constant for all Q^2 values being equal to one as one would expect. ΔH_{sea}^U while varying with Q^2 , remains small around 0.08 – 0.2 for a range of $Q^2 = 2 - 10 \text{ GeV}^2$. However $\Delta H_{\text{gluon}}^U$ is fairly large and grows rapidly with Q^2 , reaching to 4.4 at $Q^2 = 10 \text{ GeV}^2$. Such a large value, of course, is not unexpected according to QCD evolution equation. Thus, by considering only the numerical values of eq. (13), it is impossible to build a spin $\frac{1}{2}$ valon without an extra element. It turns out that the orbital angular momentum of sea partons in a valon is the extra element in the following sum rule

$$S_z^U = \frac{1}{2}(S_z^{\text{valence}} + S_z^{\text{sea}})^U + (S_z^{\text{gluon}})^U + L_z^U = \frac{1}{2} \quad (10)$$

It is known that there is a mixing operator which dresses any valence quark wave function with $\bar{q} - q$ pairs to respect PCAC. The vacuum of these massive valons is a coherent superposition of *cooper* pairs of massless quark, which resembles the structure of a valon to that of superconductivity. Studies of anisotropic superconductivity shows that a particle of condensed fluid surrounded by a cloud of correlated particles must rotate around it with the axis of rotation l . The presence of anisotropy leads to axial asymmetry of pairing correlations around the anisotropy direction. The axis of asymmetry can be associated with the polarization vector of the valence quark located at the center of the cloud. It can be shown that the magnitude of this l_z associated with the so-called *back flow* must be negative, which is precisely what we need to compensate the growth of gluon polarization contribution. In figure (5) this fact is depicted.

1. ZEUS Coll., S. Chrkanov, *et al.*, Nucl. Phys. B **637**(2002) 3.
2. F. Arash, A. N. Khorramian, Phys. Rev. C **67** (2003) 045201.
3. F. Arash, Phys. Lett. B **557** (2003) 38.
4. F. Arash, Phys. Rev. D **679** (2004) 054024.
5. J. S. Conway *et al.*, Phys. Rev. D **39**, 92 (1989).